



Time Allowed: 90 minutes

B

Max. Marks: **60**

Name: _____ ID. No. _____

Instructor's Name: _____ Section: _____ Class Time _____

Question	1	2	3	4	5	Total marks (out of 60)
Marks Obtained						

Marks out of 30

- Instructions:**
- 1) Attempt all FIVE (5) questions.
 - 2) Calculators and electronic devices are not allowed.
 - 3) Switch off your mobile(s) and put in your pocket during examination period.

Q.1 (a) If $y = \underline{\cosh}(\cos x) + \tanh^{-1}(\underline{\sqrt{x}})$, then find $\frac{dy}{dx}$. (5 marks)

$$y' = -\sinh(\cos x) \cdot \sin x + \frac{\frac{1}{2\sqrt{x}}}{1-x}$$

$$\frac{d}{x} \sqrt{x} = \frac{1}{2\sqrt{x}}$$

$$y' = -\sinh(\cos x) \cdot \sin x + \frac{1}{(1-x)2\sqrt{x}}$$

$$y' = -\sinh(\cos x) \cdot \sin x + \frac{1}{2\sqrt{x} - 2x\sqrt{x}}$$

(b) Find the limit $\lim_{x \rightarrow \infty} (x^3 e^{-x^3})$ $\infty \cdot \frac{1}{e^\infty} : \infty \cdot 0$ I.D.F (5 marks)

$$\lim_{x \rightarrow \infty} \frac{x^3}{e^{x^3}} = \frac{\infty}{\infty} \quad \text{I.D.F}$$

L'H

$$\lim_{x \rightarrow \infty} \frac{3x^2}{3x^2 e^{x^3}}$$

$$\lim_{x \rightarrow \infty} \frac{1}{e^{x^3}} = \frac{1}{\infty} = 0$$

Q.2 (a) Express the integral $\int_0^{\frac{\pi}{4}} (3 + \tan x) dx$ as a **limit of sums**. (5 marks)

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x$$

$$\Delta x = \frac{b-a}{n}$$

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n 3 + \tan \frac{\pi}{4n} i - \frac{\pi}{4n}$$

$$\Delta x = \frac{\frac{\pi}{4}}{n} = \frac{\pi}{4n}$$

$$x_i = a + \Delta x i$$

$$x_i = \frac{\pi}{4n} i$$

$$f(x_i) = 3 + \tan \frac{\pi}{4n} i$$

(b) Evaluate $\int \frac{e^{\sqrt{x}}}{\sqrt{x}} dx$

$$2 \int e^{\sqrt{x}} \cdot \frac{1}{2\sqrt{x}} dx$$

$f(x)$
 $\underbrace{\frac{1}{2\sqrt{x}}}_{f'(x)}$

$$\sqrt{x} \rightarrow \frac{1}{2\sqrt{x}}$$

$$= 2 e^{\sqrt{x}} + C$$

L.H

L.H

Q.3 (a) Evaluate $\int \frac{\sec^2 x}{\sqrt{3+\tan x}} dx$ (5 marks)

$$\begin{aligned}
 & \int \underbrace{\sec^2 x}_{f'(x)} \underbrace{(3+\tan x)^{-\frac{1}{2}}}_{f(x)} dx \\
 &= \frac{(3+\tan x)^{\frac{1}{2}}}{\frac{1}{2}} + C \\
 &= 2(3+\tan x)^{\frac{1}{2}} + C \\
 &= 2\sqrt{3+\tan x} + C
 \end{aligned}$$

(b) Find the area of the bounded region $y = x$, $y = e^x$, $0 \leq x \leq 2$.


(5 marks)

$$A: \int_0^2 e^x - x dx$$

$$A: [e^x - \frac{x^2}{2}]_0^2$$

$$A: e^2 - 2 - 1$$

$$A: e^2 - 3$$



$$y = x, \quad y = e^x$$

$$y = 1, \quad y = e$$

$$y = x < y = e^x$$

Q.4 (a) Find the **volume** of the solid obtained by rotating the bounded region

$$2x = y^2, \quad x = 0, \quad 0 \leq y \leq 2; \quad \text{about the } y\text{-axis.} \quad (5 \text{ marks})$$


$$x = \frac{y^2}{2}$$

$$V = \int_0^2 \pi [R^2 - r^2] dy$$

$$V = \pi \int_0^2 \frac{y^4}{4} dy$$

$$V = \pi \left[\frac{y^5}{20} \right]_0^2$$

$$V = \frac{32\pi}{20} = \frac{16\pi}{10} = \frac{8\pi}{5}$$



(b) Find the **volume** of the solid obtained by rotating the bounded region

$$x = -y^2 + 2y, \quad x = 0, \quad 0 \leq y \leq 2; \quad \text{about the } x\text{-axis.} \quad (5 \text{ marks})$$

(Hint: Use shell method)

$$V = \int_0^2 2\pi rh dy$$


$$V = 2\pi \int_0^2 (-y^2 + 2y)(y) dy$$

$$V = 2\pi \int_0^2 -y^3 + 2y^2 dy$$

$$V = 2\pi \left[-\frac{y^4}{4} + \frac{2y^3}{3} \right]_0^2$$

$$V = 2\pi \left[-4 + \frac{16}{3} \right]$$

$$V = 2\pi \left[\frac{4}{3} \right] = \frac{8\pi}{3}$$



$$x = -y^2 + 2y$$

$$x = 0$$

$$y = 0, \quad y = 2$$


Q.5 Choose the **correct** answer for each of the following statements.

(20 marks)

- i) If $\int_1^4 f(x) dx = 3$ and $\int_1^4 g(x) dx = 2$,
then $\int_1^4 [3f(x) - 2g(x)] dx = 5$. a) True b) False
- $\underbrace{3}_{\cdot 3} - \underbrace{2}_{\cdot 2}$
 \downarrow
 $9 - 4$
- ii) $\sinh^2 x = 1 + \cosh^2 x$ a) True b) False
- $\cosh^2 - \sinh^2 = 1$
 $\cosh^2 - \sinh^2 = -1$
- iii) $\int_1^e \frac{1}{x} dx = 1$ a) True b) False
- $\left[\ln|x| \right]_1^e$
 $1 - 0 = 1$
- iv) $\frac{d}{dx} \left[\int_{\frac{1}{x^2}}^1 e^t dt \right] = -x$ a) True b) False
- $\frac{\cos x}{\sin}$
- v) $\int \cos x \csc x dx = \frac{\cos^2 x}{2} + C$ a) True b) False
- $\int \cos x (\sin x)^{-1} : \ln |\sin x|$
- vi) $\lim_{x \rightarrow \infty} \left(\frac{1}{x} - \frac{1}{x^2} \right) = 1$ a) True b) False
- $0 - 0 = 0$
- vii) $\lim_{x \rightarrow \infty} \left(\frac{\ln x}{x} \right) = 0$ a) True b) False
- $\frac{\infty}{\infty} = \frac{\frac{1}{x}}{1} = \frac{1}{x} = 0$
- $\frac{x}{x}$
- viii) $\int \tan x dx = \ln|\sin x| + C$ a) True b) False
- ix) $\int 2x e^{(x^2+1)} dx = \frac{1}{2} e^{(x^2+1)} + C$ a) True b) False
- x) The integral that represents the volume of the solid obtained by rotating the region bounded by $y = x^2$, $x = y$, $0 \leq x \leq 1$; about the line $y = 2$ is;
 $V = \int_0^1 \pi [(2 - x^2)^2 - (2 - x)^2] dx$. a) True b) False


$$A = \pi \int_0^1 [R^2 - r^2] dx$$

$$\pi \int_0^1 (2-x^2)^2 - (2-x)^2 dx$$



$$A = 2\pi \int_0^1 rh dy$$

$$A = 2\pi \int_0^1 (\sqrt{y} - y)(2-y) dy$$



$$y = \sqrt{y}$$

$$y^2 = y$$

$$y^2 - y = 0$$

$$y(y-1) = 0$$

$$\int \frac{\alpha}{\sqrt{\alpha+x}} dx$$

$$u = x+2$$

$$du = 1 dx$$

$$\int \frac{u-2}{\sqrt{u}} du$$

$$x = u-2$$

$$\int \frac{u-2}{\sqrt{u}} du - 2 \int \frac{1}{\sqrt{u}} du$$

$$\int u^{\frac{1}{2}} du - 2 \int u^{-\frac{1}{2}} du$$

$$\frac{2}{3} u^{\frac{3}{2}} - 4 u^{\frac{1}{2}} + C$$

$$\frac{2}{3} (x+2)^{\frac{3}{2}} - 4 (x+2)^{\frac{1}{2}} + C$$

$$\int \frac{1}{4+x^2} dx = \frac{1}{2} \tan^{-1} \frac{x}{2} + C$$

$$\begin{aligned}\int \frac{3}{5+x^2} dx &= 3 \int \frac{1}{5+x^2} dx \\ &= 3 \cdot \frac{1}{\sqrt{5}} \tan^{-1} \frac{x}{\sqrt{5}} + C.\end{aligned}$$

$$\int \frac{9}{9+x^2} dx$$

$$\begin{aligned}9 \int \frac{1}{9+x^2} dx \\ &= 9 \cdot \frac{1}{3} \tan^{-1} \frac{x}{3} + C.\end{aligned}$$